To the study of the lichen genus *Usnea* (Parmeliaceae) in Kunashir Island (Sakhalin Region, Russia)

L. V. Gagarina¹, A. K. Ezhkin²

¹Komarov Botanical Institute of the Russian Academy of Sciences, St. Petersburg, Russia

²Institute of Marine Geology and Geophysics of the Far Eastern Branch of the Russian Academy of Sciences, Yuzhno-Sakhalinsk, Russia

Corresponding author: L. V. Gagarina, gagarinalv@binran.ru

Abstract. Nine species from the genus *Usnea* were found in the Kunashir Island. Six species are new for the Kunashir Island. These are *U. cornuta*, *U. glabrata*, *U. fragilescens*, *U. lapponica*, *U. pangiana*, *U. subfloridana*. In total, six secondary medullary substances detected by HPTLC were found in the species of *Usnea* from Kunashir. Four new chemotypes are described. The paper provides a key for identification of *Usnea* species in the Kunashir Island, list of specimens examined, lists of chemotypes detected by authors and known from literature.

Keywords: chemistry, distribution, key, lichen substances, Kuril Islands.

К изучению лишайников рода *Usnea* (Parmeliaceae) на острове Кунашир (Сахалинская область, Россия)

Л. В. Гагарина¹, А. К. Ежкин²

¹Ботанический институт им. В. Л. Комарова РАН, Санкт-Петербург, Россия

²Институт морской геологии и геофизики ДВО РАН, Южно-Сахалинск, Россия

Автор для переписки: Л. В. Гагарина, gagarinalv@binran.ru

Резюме. Девять видов рода *Usnea* выявлено на о. Кунашир. Шесть видов впервые обнаружены на Кунашире: *U. cornuta*, *U. glabrata*, *U. fragilescens*, *U. lapponica*, *U. pangiana*, *U. subfloridana*. Всего в талломах лишайников выявлено 6 лишайниковых веществ, идентифицировано 4 новых хемотипа. Приведен ключ для определения видов рода *Usnea* Кунашира, список изученных образцов, данные о выявленных и известных хемотипах.

Ключевые слова: ключ, лишайниковые вещества, распространение, химия, Курильские острова.

Kunashir is the southernmost island of the Kuril Islands, a volcanic archipelago in the Sea of Okhotsk in the northwest of the Pacific Ocean. The Kunashir Island with an area of 1490 square kilometers is currently administrated by the Sakhalin Region. The climate is temperate of monsoon type with heavy precipitation especially in autumn to 1400 mm and maximum temperatures in August to 18°C in average. Kunashir is included in the Sakhalin-Hokkaido province of the East Asian floristic region (Takhtadzhyan, 1978). Major part of Kunashir territory is covered by coniferous forests mostly formed by *Abies sachalinensis* (F. Schmidt) Mast., *Picea jezoensis* (Siebold et Zucc.) Carrière, and *P. glehnii* (F. Schmidt) Mast. A southern part of the is-

land is covered by broad-leaved forests mostly formed by *Quercus crispula* Blume and *Ulmus laciniata* (Trautv.) Mayr.

The lichens of Kunashir are still poorly studied. A few areas of the island has been investigated and only particular groups of lichens has been previously revised (Inсарov, Pchelkin, 1988; Titov, 1991; Bredkina *et al.*, 1992; Dombrovskaya, 1996; Tcha-banenko, 1999; Joneson *et al.*, 2004; Makryi *et al.*, 2010; Ezhkin, Kordyukov, 2016). To date, approximately 300 lichen species are known from the literature, therefore new and noteworthy records are highly expected for this territory.

The genus *Usnea* Dill. ex Adans. (Parmeliaceae, Ascomycota) consists of 355 widely distributed species (Wijayawardene *et al.*, 2020). Of these, 40 species are known from Russia (Golubkova, 1996; Spisok..., 2010; Kharpukhaeva, 2011). The genus *Usnea* contains fruticose lichens with a central axis consisting of a cartilaginous stand of longitudinal hyphae. Thallus attached by a holdfast, pale greenish or yellow green, occasionally reddish brown or blackening. Branches cylindrical, fusiform, tapering or irregular, with or without papillae, fibrils, soralia, pseudocyphellae. Apothecia rare. Cortex composed of radially or irregularly oriented hyphae. Medulla lax, dense or compact, sometimes pink pigmented. Photobiont trebouxoid. Cortex contains usnic acid, chemistry of medulla is highly variable (Halonen *et al.*, 1998; Articus, 2004; Randle *et al.*, 2009; Clerc, 2011). The delimitation of *Usnea* has been discussed by numerous authors (e. g., Ohmura, 2001, 2002, 2012; Articus, 2004; Divakar *et al.*, 2017; Ohmura, Kashiwadani, 2018) and the concept of the genus has been changed over time.

Prior our study, only three *Usnea* species were known from Kunashir Island: *U. diffracta*, *U. longissima*, and *U. rubicunda* (Bredkina *et al.*, 1992). The aim of our study is a revision of lichens of the genus *Usnea* of Kunashir.

Material and methods

Material from expeditions in Kunashir Island collected by Alexander Ezhkin in 2013–2017 (Fig. 1) was studied. Lichen identification was carried out by Ludmila Gagarina. The voucher specimens are kept in herbarium of the Institute of Marine Geology and Geophysics FEB RAS (SAK) and in herbarium of the laboratory lichenology and bryology of Komarov Botanical Institute (LE). The morphology and anatomy were studied using a light microscope and a dissecting microscope. HPTLC was performed according to the standard procedure (Culberson, Ammann, 1979; Kranner *et al.*, 2002) using solvent systems A and B by Ludmila Gagarina in the laboratory of lichenology and bryology of Komarov Botanical Institute.

Results

Nine species of the genus *Usnea* were found in Kunashir Island: *U. cornuta*, *U. diffracta*, *U. glabrata*, *U. fragilescens*, *U. lapponica*, *U. longissima*, *U. pangiana*, *U. rubicunda*, *U. subfloridana*. Of these, *U. cornuta*, *U. glabrata*, *U. fragilescens*, *U. lapponica*, *U. pangiana*, *U. subfloridana* are new for Kunashir Island.

Fig. 1. Collecting localities in Kunashir Island.

Key to the species of *Usnea* in Kunashir Island

1. Thallus pendulous, decorticated on the main branch, with numerous perpendicular fibrils *U. longissima*
- Thallus shrubby or subpendent, corticated 2
2. Branches with annular-pseudocyphellae (bamboo-like appearance), thallus isotomic-dichotomously branched *U. diffracta*
- Branches without annular-pseudocyphellae, thallus anisotomic- or isotomic-dichotomously branched 3
3. Red pigment present in cortex *U. rubicunda*
- Red pigment absent in cortex or present in medulla 4
4. Secondary branches distinctly and more or less consistently constricted at the point of attachment 5
- Secondary branches not constricted at the point of attachment 7
5. Soralia present, these generally bearing soredia, isidia always absent *U. glabrata*
- Soralia absent or, if present, then at least in part bearing isidia 6
6. Thallus clearly shrubby, upper branches unevenly anisotomically branched, readily divisible into main stems and secondary branches, isidiate soralia are concentrated at branch tips, becoming confluent *U. cornuta*

- Thallus shrubby to subpendent, upper branches evenly isotomical branched, generally not readily divisible into main stems and secondary branches, isidiate soralia are scattered over branches, remain discrete..... *U. fragilecsens*
- 7. Thallus shrubby, with distinctive concave large soralia, becoming expanded, isidia often absent (sometimes tuberculate soralia may contain spinules), medulla thick and loose *U. lapponica*
- Thallus erect to subpendent, with punctiform or convex soralia, smaller than the branch diam., isidia rare or abundant.....8
- 8. Thallus with jet black base, with small to enlarged soralia, which often contain abundant isidia..... *U. subfloridana*
- Thallus with concolor to dark brown base, with punctiform soralia, rarely with isidia. *U. pangiana*

The species

Usnea cornuta Körb. Ten chemotypes have been reported in literature for *U. cornuta*: 1) usnic, salazinic, and protocetraric ($\pm$) acids in Europe (Clerc, 1987; James *et al.*, 2009; Randlane *et al.*, 2009) and Japan (Asahina, 1956; Ohmura, 2001); 2) usnic, stictic, norstictic, and salazinic ($\pm$) acids in Europe (Clerc, 1987; James *et al.*, 2009; Randlane *et al.*, 2009); 3) usnic, lobaric, and norstictic acids in North America (Clerc, 2007); 4) usnic and norstictic acids in North America (Clerc, 2007); 5) usnic and protocetraric acids in North America (Clerc, 2007) and Japan (Asahina, 1956; Ohmura, 2001); 6) usnic, norstictic, protocetraric, and salazinic acids in Russian Far East (Ohmura *et al.*, 2017) and Taiwan (Ohmura, 2012); 7) usnic, norstictic, menegazziaic, stictic, and constictic acids in Russian Far East (Ohmura *et al.*, 2017); 8) usnic acid in North America (Clerc, 2007) and Russian Far East (Ohmura *et al.*, 2017); 9) usnic and salazinic acids in Japan (Asahina, 1956). Stevens (1999) reported occurrence of diffractaic, fumarprotocetraric, and galbinic acids in Australia.

One chemical race was found — usnic and salazinic acids.

Specimens examined: Mendelev Volcano, NW slope, 43°59'29.5"N, 145°43'20.0"E, 328 m a. s. l., spruce-fir forest, bark of *Abies sachalinensis*, 28 VII 2013, Ezhkin, SAK 324; vicinity of Mendelev Volcano, 43°59'33.1"N, 145°46'15.4"E, 161 m a. s. l., spruce-fir forest, bark of *A. sachalinensis*, 21 VI 2014, Ezhkin, SAK 337; *ibidem*, 44°00'05.5"N, 145°47'14.6"E, 37 m a. s. l., spruce-fir forest, bark of *A. sachalinensis*, 17 VI 2014, Ezhkin, SAK 349.

U. diffracta Vain. Six chemotypes have been reported for *U. diffracta*: 1) usnic, diffractaic, barbatic, baeomycesic, squamatic, and bourgeanic ($\pm$) acids, atranorin ($\pm$) in Japan (Ohmura, 2001), Taiwan (Ohmura, 2012), South Korea (Jayalal *et al.*, 2013), and Russian Far East (Ohmura *et al.*, 2017); 2) usnic and diffractaic acids in Japan (Asahina, 1956); 3) usnic, diffractaic, barbatic, 4-O-demethylbarbatic, baeomycesic, squamatic, and salazinic acids in Japan (Ohmura, 2001); 4) usnic, barbatic, 4-O-demethylbarbatic, baeomycesic, squamatic, and salazinic acids in Japan (Ohmura 2001); 5) usnic and bourgeanic acids in Japan (Ohmura, 2001); 6) usnic, barbatic, baeomycesic, and squamatic acids in South Korea (Jayalal *et al.*, 2013).

One chemical race was found in Kunashir — usnic and diffractaic acids.

Specimens examined: Mendelev Volcano surroundings, Mt. Mechnikov, 43°58'42.0"N, 145°42'52.1"E, 340 m a. s. l., dark coniferous forest, bark of *Picea glehnii*, 6 VIII 2013, Ezhkin, SAK

1257; *ibidem*, 44°00'15.3"N, 145°41'53.5"E, 201 m a. s. l., mixed forest, bark of *Quercus crispula*, 1 VIII 2013, *Ezhkin*, SAK 1256; Mendelev Volcano, NW slope, 43°59'58.8"N, 145°42'46.5"E, 343 m a. s. l., dark coniferous forest, bark of *Picea glehnii*, 5 VIII 2015, *Ezhkin*, SAK 1255; Dubovoye village surroundings, 43°47'47.0"N, 145°30'15.3"E, 107 m a. s. l., dark coniferous forest, bark of *Quercus crispula*, 23 VIII 2017, *Ezhkin*, SAK 1258; Belkina River valley, 43°54'10.9"N, 145°35'51.1"E, 56 m a. s. l., old-growth mixed coniferous-broadleaved forest, bark of *Ulmus laciniata*, 23 VI 2014, *Ezhkin*, SAK 1254; Rurui Volcano surroundings, Prosolovskie springs, 44°19'58.4"N, 146°01'14.5"E, 16 m a. s. l., old-growth mixed coniferous-broadleaved forest, bark of *Sorbus commixta* Hedl., 19 VI 2014, *Ezhkin*, SAK 1259; Tretyakova village surroundings, 43°53'23.5"N, 145°36'37.7"E, 37 m a. s. l., old-growth mixed coniferous-broadleaved forest, bark of *Ulmus laciniata*, 23 VI 2014, *Ezhkin*, SAK 1253.

Usnea fragilesceus Hav. ex Lynge. Seven chemotypes have been reported in literature for *U. fragilesceus*: 1) usnic, norstictic, salazinic, and protocetraric ($\pm$) acids in Russian Far East (Ohmura *et al.*, 2017); 2) usnic, norstictic, stictic ($\pm$), menegazziaic ($\pm$), and fatty ($\pm$) acids in Russian Far East (Ohmura *et al.*, 2017), Japan, and Taiwan (Ohmura, 2001); 3) usnic, stictic, cryptostictic ($\pm$), and menegazziaic ($\pm$) acids in Japan and Taiwan (Ohmura, 2001); 4) usnic, salazinic, stictic acid complex ($\pm$), atranorin ($\pm$) in Japan and Taiwan (Ohmura, 2001); 5) usnic, psoromic, and 2-O-demethylpsoromic acids in Japan and Taiwan (Ohmura, 2001); 6) usnic acid in Japan and Taiwan (Ohmura, 2001); 7) usnic, stictic acid complex in Europe (Randlane *et al.*, 2009); 8) usnic, salazinic acids in North America (Clerc, 2007); 9) usnic, psoromic acids in North America (Clerc, 2007).

Two chemotypes were found — usnic, salazinic acids; usnic acid.

Specimens examined: Belkina River valley, 43°54'12.1"N, 145°36'4.1"E, 77 m a. s. l., mixed coniferous-deciduous forest, bark of *Betula ermanii* Cham., 23 VI 2014, *Ezhkin*, SAK 341, 343 (SAK 341 — without medulla substances, SAK 343 — salazinic acid); Mendelev Volcano surroundings, Mechnikov Mt., 43°58'33.7"N, 145°43'19.8"E, 469 m a. s. l., spruce-fir forest, bark of *Picea jezoensis*, 6 VIII 2013, *Ezhkin*, SAK 351.

U. glabrata (Ach.) Vain. Six chemotypes have been reported in literature for *U. glabrata*: 1) usnic, protocetraric, and fumarprotocetraric acids in Europe and North America (Randlane *et al.*, 2009); 2) usnic, salazinic, and norstictic acids in Fennoscandia (Halonen *et al.*, 1999); 3) usnic acid in Europe and North America (Clerc, 1987; Myllys, 1994; Brodo *et al.*, 2001); 4) usnic, salazinic acids in Japan (Asahina, 1956) and Russian Far East (Ohmura *et al.*, 2017); 5) usnic, barbatic, 4-O-demethylbarbatic (trace), norstictic ($\pm$), and protocetraric ($\pm$) acids in Russian Far East (Ohmura *et al.*, 2017); 6) usnic acid in Europe (Brodo *et al.*, 2001).

Four chemotypes were found — usnic, protocetraric, salazinic, thamnolic acids; usnic, salazinic acids; usnic, protocetraric, salazinic acids; usnic acid. Chemotype with thamnolic acid was not previously known for this species. Chemotype with usnic, protocetraric, and salazinic acids together is new for Kunashir.

Specimens examined: Belkina River valley, 43°53'24.8"N, 145°36'28.3"E, 66 m a. s. l., mixed coniferous-deciduous forest, bark of *Abies sachalinensis*, 23 VI 2014, *Ezhkin*, SAK 363; *ibidem*, 43°53'21.9"N, 145°36'54.7"E, 30 m a. s. l., mixed coniferous-deciduous forest, bark of *Betula ermanii*, 22 VI 2014, *Ezhkin*, SAK 371, 367, 369; *ibidem*, 43°54'12.1"N, 145°36'04.1"E, 77 m a. s. l., mixed coniferous-deciduous forest, bark of *B. ermanii*, 23 VI 2014, *Ezhkin*, SAK 344; vicinity of Lagunnoe Lake,

44°02'49.7"N, 145°45'59.1"E, 65 m a. s. l., mixed coniferous-deciduous forest, bark of *Ulmus laciniata*, 17 VIII 2013, *Ezhkin*, SAK 332; vicinity of Mendelev Volcano, 44°00'05.9"N, 145°47'17.6"E, 29 m a. s. l., spruce-fir forest, bark of *Abies sachalinensis*, 17 VI 2014, *Ezhkin*, SAK 352.

U. lapponica Vain. Three chemotypes have been reported in literature for *U. lapponica*: 1) usnic, salazinic acids with different medullary substances (protocetraric, barbatic, caperatic acids, zeorin) in Europe, North America (Halonen *et al.*, 1998), and Russian Far East (Ohmura *et al.*, 2017); 2) usnic and psoromic acids in Europe (Halonen *et al.*, 1999); 3) usnic acid in Europe and North America (Randlane *et al.*, 2009).

Three chemotypes were found — usnic, salazinic, barbatic acids; usnic acid; usnic, salazinic acids.

Specimens examined: Aliger Lake surroundings, 44°01'45.8"N, 145°43'12.7"E, 87 m a. s. l., mixed coniferous-deciduous forest, bark of *Betula ermanii*, 20 VI 2014, *Ezhkin*, SAK 318; vicinity of Tretakova village, 43°58'28.4"N, 145°39'49.3"E, 163 m a. s. l., mixed coniferous-deciduous forest, bark of *Ulmus laciniata*, 22 VI 2014, *Ezhkin*, SAK 321; *ibidem*, 43°53'23.5"N, 145°36'37.7"E, 36 m a. s. l., old-growth mixed coniferous-broadleaved forest, bark of *U. laciniata*, 23 VI 2014, *Ezhkin*, SAK 1240; Mendelev Volcano surroundings, 43°59'30.4"N, 145°45'59.7"E, 183 m a. s. l., spruce-fir forest, bark of *Abies sachalinensis*, 24 VI 2014, *Ezhkin*, SAK 335.

U. longissima Ach. Ten chemotypes have been reported for *U. longissima*: 1) usnic and barbatic acids in Japan (Ohmura, 2001), Europe (Randlane *et al.*, 2009), North America (Halonen *et al.*, 1998); 2) usnic and diffractaic acids in Japan (Ohmura, 2001), Europe (Randlane *et al.*, 2009), North America (Halonen *et al.*, 1998); 3) usnic and evernic acids in Japan (Ohmura, 2001), Europe (Randlane *et al.*, 2009), North America (Halonen *et al.*, 1998), and Russian Far East (Ohmura *et al.*, 2017); 4) usnic and salazinic acids in Japan (Asahina, 1956); 5) usnic and fumarprotocetraric acids in Japan (Asahina, 1956); 6) usnic, barbatic, 4-O-demethylbarbatic, and squamatic ($\pm$) acids in Taiwan (Ohmura, 2012); 7) usnic, diffractaic, and barbatic acids in Taiwan (Ohmura, 2012); 8) usnic, evernic, and lecanoric acids in Taiwan (Ohmura, 2012); 9) usnic, barbatic (trace), diffractaic, baeomycesic, and squamatic acids in Russian Far East (Ohmura *et al.*, 2017); 10) usnic acid in Europe and North America (Halonen, 1998).

One chemical race was recognized in the study area — usnic acid.

Specimens examined: Zmeinyi Creek valley, 44°00'27.6"N, 145°41'20.8"E, 99 m a. s. l., mixed coniferous-deciduous forest, bark of *Hydrangea paniculata* Siebold, 31 VII 2013, *Ezhkin*, SAK 357; *ibidem*, 44°00'28.1"N, 145°41'39.0"E, 158 m a. s. l., mixed coniferous-deciduous forest, bark of *Abies sachalinensis*, 31 VII 2013, *Ezhkin*, SAK 340; Belkina River valley, 43°54'10.9"N, 145°35'51.1"E, 56 m a. s. l., mixed coniferous-deciduous forest, bark of *Ulmus laciniata*, 23 VI 2014, *Ezhkin*, SAK 348; *ibidem*, 43°53'51.5"N, 145°36'22.5"E, 70 m a. s. l., mixed coniferous-deciduous forest, bark of *Abies sachalinensis*, 23 VI 2014, *Ezhkin*, SAK 339.

U. pangiana Stirt. Three chemotypes have been reported for *U. pangiana*: 1) usnic, salazinic, norstictic ($\pm$), protocetraric ($\pm$), constictic ($\pm$), and squamatic ($\pm$) acids in Taiwan, Japan (Ohmura, 2001, 2012), and Russian Far East (Ohmura *et al.*, 2017); 2) usnic, salazinic, barbatic, 4-O-demethylbarbatic, norstictic ($\pm$), and protocetraric

(±) acids, atranorin (±) in Taiwan and Japan (Ohmura, 2001, 2012); 3) usnic acid in Russian Far East (Ohmura *et al.*, 2017).

Two chemical races were found — usnic, salazinic acids; usnic, salazinic, barbatic acids.

Specimens examined: Belkina River valley, 43°53'21.9"N, 145°36'54.7"E, 30 m a. s. l., mixed coniferous-deciduous forest, bark of *Betula ermanii*, 22 VI 2014, *Ezhkin*, SAK 373, 372, 370, 368; *ibidem*, 43°54'12.1"N, 145°36'04.1"E, 77 m a. s. l., mixed coniferous-deciduous forest, bark of *B. ermanii*, 23 VI 2014, *Ezhkin*, SAK 345, 342; *ibidem*, 43°54'13.1"N, 145°36'15.4"E, 57 m a. s. l., mixed forest, bark of *Salix caprea* L., 23 VI 2014, *Ezhkin*, SAK 338; *ibidem*, 43°53'23.5"N, 145°36'37.7"E, 36 m a. s. l., mixed coniferous-deciduous forest, bark of *Betula ermanii*, 23 VI 2014, *Ezhkin*, SAK 374; vicinity of Tretiakova village, 43°58'34.1"N, 145°39'45.4"E, 191 m a. s. l., mixed coniferous-deciduous forest, bark of *Taxus cuspidata* Siebold et Zucc. ex Endl., 22 VI 2014, *Ezhkin*, SAK 319; *ibidem*, 43°58'28.4"N, 145°39'49.3"E, 163 m a. s. l., mixed coniferous-deciduous forest, bark of *Ulmus laciniata*, 22 VI 2014, *Ezhkin*, SAK 320; *ibidem*, bark of *Phellodendron sakhalinense* (F. Schmidt) Sarg., 22 VI 2015, *Ezhkin*, SAK 1246; *ibidem*, old-growth mixed coniferous-broadleaved forest, bark of *Abies sakhalinensis*, 22 VI 2015, *Ezhkin*, SAK 1247; Mendelev Volcano surroundings, Mechnikov Mt., 43°58'15.9"N, 145°43'34.1"E, 547 m a. s. l., fir forest, bark of *A. sachalinensis*, 13 VIII 2013, *Ezhkin*, SAK 333, 334; *ibidem*, 43°58'33.7"N, 145°43'19.8"E, 468 m a. s. l., coniferous forest, bark of *Sorbus commixta*, 6 VIII 2013, *Ezhkin*, SAK 1237; *ibidem*, 43°59'23.3"N, 145°47'5.1"E, 94 m a. s. l., spruce-fir forest, bark of *Abies sachalinensis*, 24 VI 2014, *Ezhkin*, SAK 336, 1248; Mendelev Volcano, 44°00'04.8"N, 145°42'26.8"E, 134 m a. s. l., mixed coniferous-deciduous forest, bark of *Picea jezoensis*, 26 VIII 2013, *Ezhkin*, SAK 326; *ibidem*, 43°59'21.8"N, 145°46'16.7"E, 197 m a. s. l., spruce-fir forest, bark of *Sorbus commixta*, 11 VIII 2015, *Ezhkin*, SAK 330; *ibidem*, 43°59'31.9"N, 145°43'15.6"E, 317 m a. s. l., coniferous forest, bark of *Abies sakhalinensis*, 28 VII 2013, *Ezhkin*, SAK 1252; Dubovoye village surroundings, 43°46'28.7"N, 145°31'00.9"E, 58 m a. s. l., oak forest, bark of *Quercus crispula*, 23 VIII 2017, *Ezhkin*, SAK 1251; *ibidem*, 43°46'47.6"N, 145°29'48.7"E, 69 m a. s. l., oak forest, bark of *Q. crispula*, 23 VIII 2017, *Ezhkin*, SAK 1239, 1238; Golovnino Volcano surroundings, 43°52'51.6"N, 145°36'20.3"E, 89 m a. s. l., mixed forest, bark of *Sorbus commixta*, 22 VIII 2017, *Ezhkin*, SAK 1242; *ibidem*, 43°52'20.9"N, 145°36'00.9"E, 89 m a. s. l., mixed forest, bark of *Quercus crispula*, 23 VIII 2017, *Ezhkin*, SAK 1241.

Usnea rubicunda Stirt. Four chemotypes have been reported in literature for *U. rubicunda*: 1) usnic and stictic acids as a main substances, and norstictic and psoromic acids as accessories in Europe (James *et al.*, 2009; Randlane *et al.*, 2009) and Japan (Asahina, 1956); 2) usnic and salazinic acids as a main substances and norstictic acid as an accessory in Europe (James *et al.*, 2009; Randlane *et al.*, 2009); 3) usnic, norstictic, stictic, menegazziaic, constictic, and squamatic (±) acids and atranorin (±) in Taiwan (Ohmura, 2012); 4) usnic, norstictic, stictic, menegazziaic, constictic, psoromic, and 2-O-demethylpsoromic (±) acids in Taiwan (Ohmura, 2012) and Russian Far East (Ohmura *et al.*, 2017).

Two chemical races were found — usnic, salazinic, norstictic acids; usnic acid. Chemotype without medullary substances (usnic acid in cortex) was not previously known for this species.

Specimens examined: Belkina River valley, 43°53'23.5"N, 145°36'37.7"E, 36 m a. s. l., mixed coniferous-deciduous forest, bark of *Abies sachalinensis*, 23 VI 2014, *Ezhkin*, SAK 360, 361, 358, 362, 359; *ibidem*, 43°53'24.8"N, 145°36'28.3"E, 66 m a. s. l., mixed coniferous-deciduous forest, bark of

A. sachalinensis, 23 VI 2014, *Ezhkin*, SAK 364, 365; Golovnino Volcano surroundings, 43°52'51.6"N, 145°36'20.3"E, 89 m a. s. l., mixed forest, bark of *A. sakhalinensis*, 22 VIII 2017, *Ezhkin*, SAK 1249, 1243.

U. subfloridana Stirt. Four chemotypes have been reported for *U. subfloridana*: 1) usnic and squamatic acids in Europe, North America (Randlane *et al.*, 2009), Russian Far East (Ohmura *et al.*, 2017), Taiwan, and Japan (Ohmura, 2001); 2) usnic and thamnolic acids in Europe, North America (Randlane *et al.*, 2009), Taiwan (Ohmura, 2012), and Japan (Ohmura, 2001); 3) usnic, squamatic, thamnolic, and alecatorialic ($\pm$) acids in Europe, North America (Randlane *et al.*, 2009), Taiwan, and Japan (Ohmura, 2001); 4) usnic, norstictic acids in Taiwan and Japan (Ohmura, 2001).

One chemotype was found — usnic, salazinic, barbatic acids. Chemotype with salazinic and barbatic acids was not previously known for this species. Among known *Usnea* species, *U. wasmuthii* Räsänen has similar chemotype, but differs by its morphological features, e. g. by more excavate soralia with few or without isidiomorphs and basal part with tiny, lenticular, longitudinally oriented cracks. All studied specimens lacked mentioned features and had typical morphology for *U. subfloridana*.

Specimens examined: Belkina River valley, 43°53'51.5"N, 145°36'22.5"E, 70 m a. s. l., mixed coniferous-deciduous forest, bark of *Sorbus commixta*, 23 VI 2014, *Ezhkin*, SAK 346; Dubovoye village surroundings, 43°46'28.7"N, 145°31'00.9"E, 58 m a. s. l., oak forest, bark of *Quercus crispula*, 23 VIII 2017, *Ezhkin*, SAK 1250; *ibidem*, 43°47'47.0"N, 145°30'15.3"E, 107 m a. s. l., oak forest, bark of *Q. crispula*, 23 VIII 2017, *Ezhkin*, SAK 1245; *ibidem*, 43°46'47.6"N, 145°29'48.7"E, 69 m a. s. l., oak forest, bark of *Q. crispula*, 23 VIII 2017, *Ezhkin*, SAK 1238.

Conclusion

In total, six secondary medullary substances were detected in *Usnea* species growing on Kunashir Island: barbatic, diffractaic, norstictic, protocetraric, salazinic, and thamnolic acids. Four new chemotypes were discovered: 1) *U. glabrata* — usnic, protocetraric, salazinic, thamnolic acids; 2) *U. glabrata* — usnic, protocetraric, salazinic acids; 3) *U. rubicunda* — usnic acid; 4) *U. subfloridana* — usnic, salazinic, barbatic acids. In our opinion, new chemotypes can be described from different countries as most of *Usnea* species have a relatively complex chemistry.

Acknowledgments

The work was partially supported by the Russian Foundation of Basic Researches (No. 18-04-00098 A, 19-54-50010) and institutional research project "Flora and systematics of lichens and mosses in Russia and phytogeographically important regions" (AAAA-A19-119020690077-4). We are grateful to Yoshihito Ohmura (National Museum of Nature and Science, Japan) for identification of *U. pangiana* sample with using DNA methods. We thank Alexander Kordyukov (Institute of Marine Geology and Geophysics FEB RAS, Yuzhno-Sakhalinsk, Russia) for preparing the map. The authors also thank Andrei Tsurykau (Francisk Skorina Gomel State University, Gomel, Belarus; Institute of Natural Sciences, Samara National Research University, Samara, Russia) for valuable comments on the manuscript.

References

- Articus K. 2004. *Neuropogon* and the phylogeny of *Usnea* s. l. (Parmeliaceae, Lichenized Ascomycetes). *Taxon* 53(4): 925–934. <https://doi.org/10.2307/4135560>
- Asahina Y. 1956. *Lichens of Japan. Vol. 3. Genus Usnea*. Tokyo: 129 p.
- Bredkina I. I., Dobrysh A. A., Makarova I. I., Titov A. N. 1992. To the lichen flora of Kunashir Island (Kuril Islands). *Novosti sistematiki nizshikh rastenii* 28: 90–94. (In Russ.).
- Brodo I. M., Sharnoff D. S., Sharnoff S. 2001. *Lichens of North America*. New Haven: 795 p.
- Clerc P. 1987. On the morphology of soralia in the genus *Usnea*. *Bibliotheca lichenologica* 25: 99–107.
- Clerc P. 2007. *Usnea. Lichen flora of the Greater Sonoran Desert Region. Vol. 3*. Tempe: 302–335.
- Clerc P. 2011. *Usnea. Nordic lichen flora. Vol. 4*. Uppsala: 107–127.
- Culberson C. F., Ammann K. K. 1979. Standardmethode zur Dünnschichtchromatographie von Flechtensubstanzen. *Herzogia* 5: 1–24. (In Germ.).
- Divakar P. K., Crespo A., Kraichak E., Leavitt S. D., Singh G., Schmitt I., Lumbsch H. T. 2017. Using a temporal phylogenetic method to harmonize family- and genus-level classification in the largest clade of lichen-forming fungi. *Fungal Diversity* 84(1): 101–117. <https://doi.org/10.1007/s13225-017-0379-z>
- Dombrovskaya A. V. 1996. *Rod Stereocaulon na territorii byvshego SSSR* [Genus *Stereocaulon* in the former USSR]. St. Petersburg: 270 p. (In Russ.).
- Ezhkin A. K., Kordyukov A. V. 2016. Features of changes in the parameters of the epiphytic lichen cover in the vicinity of the Mendeleyev volcano (Kunashir Island, South Kuriles). *Bulletin of the Botanical Garden and Institute FEB RAS* 15: 26–28. (In Russ.).
- Golubkova N. S. 1996. *Usnea. Opredelitel' lichaynikov Rossii. Vyp. 3* [Handbook of the lichens of Russia. Iss. 6]. St. Petersburg: 62–107. (In Russ.).
- Halonen P., Clerc P., Goward T., Brodo I., Wulff K. 1998. Synopsis of the genus *Usnea* (lichenized Ascomycetes) in British Columbia, Canada. *The Bryologist* 101(1): 36–60. <https://doi.org/10.2307/3244073>
- Halonen P., Myllus L., Ahti T., Petrova V. O. 1999. The lichen genus *Usnea* in East Fennoscandia. III. The shrubby species. *Annales Botanici Fennici* 36: 235–256.
- Insarov G. E., Pchelkin A. V. 1988. *Kolichestvennye kharakteristiki sostoyaniya epifitnoi likhenoflory Kuril'skogo zapovednika* [Quantitative characteristics of epiphytic lichen flora of Kuril'skiy Reserve]. Moscow: 174 p. (In Russ.).
- James P. W., Clerc P., Purvis O. W. 2009. *Usnea* Dill. ex Adans. *The Lichens of Great Britain and Ireland*. London: 918–929.
- Jayalal U., Joshi S., Oh S.-O., Jin Koh Y., Crişan F., Hur J.-S. 2013. Lichen mycota in South Korea: The Genus *Usnea*. *Mycobiology* 41(3): 126–130. <https://doi.org/10.5941/MYCO.2013.41.3.126>
- Joneson S., Kashiwadani H., Tchabanenko S., Gage S. 2004. *Ramalina* of the Kuril Islands. *The Bryologist* 107(1): 98–106. [https://doi.org/10.1639/0007-2745\(2004\)107\[98:ROTKI\]2.0.CO;2](https://doi.org/10.1639/0007-2745(2004)107[98:ROTKI]2.0.CO;2)
- Kharpukhaeva T. M. 2011. Species of the genus *Usnea* (Parmeliaceae) in Republic of Buryatia. *Novosti sistematiki nizshikh rastenii* 45: 257–266 (In Russ. with Engl. abstract). <https://doi.org/10.31111/nsnr/2011.45.257>
- Kranner I., Beckett R. P., Varma A. K. 2002. *Protocols in lichenology*. Berlin, Heidelberg: 580 p. <https://doi.org/10.1007/978-3-642-56359-1>
- Makryi T. V., Taran T. V., Tchabanenko S. I. 2010. *Collema complanatum* (Collemataceae, Lichenes) — a new species of lichen flora of Russia. *Botanicheskii Zhurnal* 95 (7): 989–991. (In Russ. with Engl. abstract).
- Myllus L. 1994. *Usnea glabrata* (lichenized Ascomycotina) in East Fennoscandia. *Acta Botanica Fennica* 150: 125–130.

- Ohmura Y. 2001. Taxonomic study of the genus *Usnea* (lichenized Ascomycetes) in Japan and Taiwan. *The Journal of the Hattori Botanical Laboratory* 90: 1–96. https://doi.org/10.18968/jhbl.90.0_1
- Ohmura Y. 2002. Phylogenetic evaluation of infrageneric groups of the genus *Usnea* based on ITS regions in rDNA. *The Journal of the Hattori Botanical Laboratory* 92: 231–243. https://doi.org/10.18968/jhbl.92.0_231
- Ohmura Y. 2012. A Synopsis of the lichen genus *Usnea* (Parmeliaceae, Ascomycota) in Taiwan. *Memoirs of National Museum of Nature and Science, Tokyo* 48: 91–137.
- Ohmura Y., Skirina I., Skirin F. 2017. Contribution to the knowledge of the genus *Usnea* (Parmeliaceae, Ascomycota) in Southern Far East Russia. *Bulletin of the National Museum of Nature and Science, Series B (Botany), Tokyo* 43(1): 1–10.
- Ohmura Y., Kashiwadani H. 2018. Checklist of lichens and allied fungi of Japan. *National Museum of Nature and Science Monographs* 49: 1–140.
- Randlane T., Tõrra T., Saag A., Saag L. 2009. Key to European *Usnea* species. *Bibliotheca lichenologica* 100: 419–462.
- Spisok lichenoflory Rossii* [A checklist of the lichen flora of Russia]. 2010. St. Petersburg: 194 p. (In Russ. with Engl. introduction).
- Stevens G. N. 1999. A revision of the lichen family Usneaceae in Australia. *Bibliotheca lichenologica* 72: 1–128.
- Takhtadzhyan A. L. 1978. *Floristicheskie oblasti Zemli* [The floristic regions of the World]. Leningrad: 248 p. (In Russ.).
- Tchabanenko S. I. 1999. Lichens of Kuril Reserve (the Kunashir Island). *Issledovanie rastitel'nogo pokrova rossiiskogo Dal'nego Vostoka. Trudy botanicheskikh sadov DVO RAN. T. 1.* [Study of the vegetation cover of the Russian Far East. Proceedings of the Botanical Gardens of the Far-East Branch of the Russian Academy of Sciences. Vol. 1.]. Vladivostok: 221–228. (In Russ.).
- Titov A. N. 1991. Lichens of the order Caliciales from the Kunashir Island (the Kuril Islands). *Botanicheskii zhurnal* 76(4): 599–601. (In Russ.).
- Wijayawardene N. N., Hyde K. D., Al-Ani L. K. T., Tedersoo L., Haelewaters D., Rajeshkumar K. C., Zhao R. L., Aptroot A., Leontyev D. V., Saxena R. K. *et al.* 2020. Outline of fungi and fungus-like taxa. *Mycosphere* 11(1): 1060–1456. <https://doi.org/10.5943/mycosphere/11/1/8>

Литература

- Articus K. 2004. *Neuropogon* and the phylogeny of *Usnea* s. l. (Parmeliaceae, Lichenized Ascomycetes). *Taxon* 53(4): 925–934. <https://doi.org/10.2307/4135560>
- Asahina Y. 1956. *Lichens of Japan. Vol. 3. Genus Usnea*. Tokyo: 129 p.
- [Bredkina *et al.*] Бредкина Л. И., Добрыш А. А., Макарова И. И., Титов А. Н. 1992. К флоре лишайников острова Кунашир (Курильские острова). *Новости систематики низших растений* 28: 90–94.
- Brodo I. M., Sharnoff D. S., Sharnoff S. 2001. *Lichens of North America*. New Haven: 795 p.
- Clerc P. 1987. On the morphology of soralia in the genus *Usnea*. *Bibliotheca lichenologica* 25: 99–107.
- Clerc P. 2007. *Usnea. Lichen flora of the Greater Sonoran Desert Region. Vol. 3*. Tempe: 302–335.
- Clerc P. 2011. *Usnea. Nordic lichen flora. Vol. 4*. Uppsala: 107–127.
- Culberson C. F., Ammann K. K. 1979. Standardmethode zur Dünnschichtchromatographie von Flechtensubstanzen. *Herzogia* 5: 1–24.
- Divakar P. K., Crespo A., Kraichak E., Leavitt S. D., Singh G., Schmitt I., Lumbsch H. T. 2017. Using a temporal phylogenetic method to harmonize family- and genus-level classification in the largest clade of lichen-forming fungi. *Fungal Diversity* 84(1): 101–117. <https://doi.org/10.1007/s13225-017-0379-z>

- [Dombrovskaya] Домбровская А. В. 1996. *Под Stereocaulon на территории бывшего СССР*. СПб.: 270 с.
- [Ezhkin, Kordyukov] Ежкин А. К., Кордюков А. В. 2016. Особенности изменения параметров эпифитного лишайникового покрова в окрестностях вулкана Менделеева (о-в Кунашир, Южные Курилы). *Бюллетень Ботанического сада-института ДВО РАН* 15: 23–25.
- [Golubkova] Голубкова Н. С. 1996. *Usnea. Определитель лишайников России. Вып. 6*. СПб.: 62–107.
- Halonen P., Clerc P., Goward T., Brodo I., Wulff K. 1998. Synopsis of the Genus *Usnea* (Lichenized Ascomycetes) in British Columbia, Canada. *The Bryologist* 101(1): 36–60.
<https://doi.org/10.2307/3244073>
- Halonen P., Myllus L., Ahti T., Petrova V. O. 1999. The lichen genus *Usnea* in East Fennoscandia. III. The shrubby species. *Annales Botanici Fennici* 36: 235–256.
- [Insarov, Pchelkin] Инсаров Г. Э., Пчелкин А. В. 1984. *Количественные характеристики состояния эпифитной лишайнофлоры Курильского заповедника*. М.: 174 с.
- James P. W., Clerc P., Purvis O. W. 2009. *Usnea* Dill. ex Adans. *The Lichens of Great Britain and Ireland*. London: 918–929.
- Jayalal U., Joshi S., Oh S.-O., Jin Koh Y., Crişan F., Hur J.-S. 2013. Lichen mycota in South Korea: The genus *Usnea*. *Mycobiology* 41(3): 126–130. <https://doi.org/10.5941/MYCO.2013.41.3.126>
- Joneson S., Kashiwadani H., Tchabanenko S., Gage S. 2004. *Ramalina* of the Kuril Islands. *The Bryologist* 107(1): 98–106. [https://doi.org/10.1639/0007-2745\(2004\)107\[98:ROTKI\]2.0.CO;2](https://doi.org/10.1639/0007-2745(2004)107[98:ROTKI]2.0.CO;2)
- [Kharpukhaeva] Харпухаева Т. М. 2011. Виды рода *Usnea* (Parmeliaceae) в Республике Бурятия. *Новости систематики низших растений* 45: 257–266.
<https://doi.org/10.31111/nsnr/2011.45.257>
- Kranner I., Beckett R. P., Varma A. K. 2002. *Protocols in lichenology*. Berlin, Heidelberg: 580 p.
<https://doi.org/10.1007/978-3-642-56359-1>
- [Makryi et al.] Макрый Т. В., Таран А. А., Чабаненко С. И. 2010. *Collema complanatum* (Collema-taceae) новый вид для лишайнофлоры России. *Ботанический журнал* 95(7): 989–991.
- Myllus L. 1994. *Usnea glabrata* (lichenized Ascomycotina) in East Fennoscandia. *Acta Botanica Fennica* 150: 125–130.
- Ohmura Y. 2001. Taxonomic study of the genus *Usnea* (lichenized Ascomycetes) in Japan and Taiwan. *The Journal of the Hattori Botanical Laboratory* 90: 1–96. https://doi.org/10.18968/jhbl.90.0_1
- Ohmura Y. 2002. Phylogenetic evaluation of infrageneric groups of the genus *Usnea* based on ITS regions in rDNA. *The Journal of the Hattori Botanical Laboratory* 92: 231–243.
https://doi.org/10.18968/jhbl.92.0_231
- Ohmura Y. 2012. A synopsis of the lichen genus *Usnea* (Parmeliaceae, Ascomycota) in Taiwan. *Memoirs of National Museum of Nature and Science, Tokyo* 48: 91–137.
- Ohmura Y., Skirina I., Skirin F. 2017. Contribution to the knowledge of the genus *Usnea* (Parmeliaceae, Ascomycota) in Southern Far East Russia. *Bulletin of the National Museum of Nature and Science, Series B (Botany), Tokyo* 43(1): 1–10.
- Ohmura Y., Kashiwadani H. 2018. Checklist of lichens and allied fungi of Japan. *National Museum of Nature and Science Monographs* 49: 1–140.
- Randlane T., Tõrra T., Saag A., Saag L. 2009. Key to European *Usnea* species. *Bibliotheca lichenologica* 100: 419–462.
- [Spisok...] *Список лишайнофлоры России*. 2010. СПб.: 194 с.
- Stevens G. N. 1999. A revision of the lichen family Usneaceae in Australia. *Bibliotheca lichenologica* 72: 1–128.
- [Takhtadzhyan] Тахтаджян А. Л. 1978. *Флористические области Земли*. Л.: 248 с.

- [Tchabanenko] Чабаненко С. И. 1999. Лишайники Курильского заповедника (остров Кунашир). *Исследование растительного покрова российского Дальнего Востока. Труды ботанических садов ДВО РАН. Т. 1.* Владивосток: 221–228.
- [Titov] Титов А. Н. 1991. Лишайники порядка Caliciales с острова Кунашир (Курильские острова). *Ботанический журнал* 76(4): 599–601.
- Wijayawardene N. N., Hyde K. D., Al-Ani L. K. T., Tedersoo L., Haelewaters D., Rajeshkumar K. C., Zhao R. L., Aptroot A., Leontyev D. V., Saxena R. K. *et al.* 2020. Outline of fungi and fungus-like taxa. *Mycosphere* 11(1): 1060–1456. <https://doi.org/10.5943/mycosphere/11/1/8>